

IMMUCOX[®]T

ORAL COCCIDIOSIS VACCINE
Live Oocysts of Eimeria spp.

APPROVED SPECIES

- Turkeys

VACCINE TYPE

- Live, sporulated Eimeria oocysts
 - Eimeria adenoeides
 - Eimeria melegrimitis

DISEASE PROTECTION

- Coccidiosis caused by the species contained in the vaccine

VACCINE PRESENTATION

- Live sporulated oocysts in a liquid suspension

ROUTE OF ADMINISTRATION

Droplet application

AGE OF ADMINISTRATION

- 1-3 Day old poults

PACKAGING (PRODUCT CODE)

- 10 x 1,000 doses (V55210C)
- Each vial contains 10 ml liquid

QUALITY

Thoroughly tested for purity, sterility, potency, and safety and compliant with all applicable USDA, CFIA and Ceva standards.

KEY ADVANTAGES

- Effective vaccination against coccidiosis caused by the 2 Eimeria species economically important to turkeys
- Gel droplet application keeps oocysts in suspension during application, preventing sedimentation of oocysts and facilitating a uniform vaccine uptake of all the species included in the vaccine
- Offer a natural cost-effective alternative to anticoccidials

See reverse side and package label for complete use directions and information.



IMMUCOX[®]T

ORAL COCCIDIOSIS VACCINE
Live Oocysts of Eimeria spp.

INDICATIONS

For the vaccination of turkeys against Coccidiosis. For more information regarding safety data, see productdata.aphis.usda.gov.

PRECAUTIONS

- Be sure that no anticoccidial (coccidiostat) is used in the feed. Even a low level of anticoccidial can reduce or negate the effectiveness of the vaccine.
- Broad spectrum antibiotics which possess anticoccidial activity (eg. Tetracyclines, sulfa drugs and biofurans) should not be used together with the vaccine for the first 16 days after vaccination.
- Do not over dose or under dose. If birds are housed in confined space (e.g. brooding rings), they must be released from confinement by the sixth day post vaccination or the area must be at least doubled.
- Vaccinate only healthy turkeys.

DIRECTIONS FOR USE

Follow the directions carefully.

For 10,000 doses add 70g of Cevagel[®] to 2.5L of cold (2-8°C) distilled water. Mix well with a blender. No further agitation is required for use on the same day. Mix the vaccine vial well prior to opening. Pour half of the gel into a clean container, add the vaccine, then add the remaining gel to a final volume of 2.5 L for 10,000 doses. Mix with a blender. Vaccinate at day of age or older with a ratio of 25mL for every 100 turkeys using a gel administration machine.

WARNING

Store vaccine at 2-8°C. DO NOT FREEZE. Do not vaccinate within 21 days of slaughter. Do not mix with other products, except as specified on this label. Contact a physician for human exposure. Inactivate unused product. Use entire contents when first opened. FOR USE IN ANIMALS ONLY.

PRESERVATIVE

Potassium Dichromate

CAUTION

Efficacy/potency in progress. Conditions of storage and handling of the vaccine by the user, administration of the vaccine, the responsiveness of individual turkeys to the vaccine and whether the directions for use and precautionary measures are being followed carefully will affect the capacity of the vaccine.

The capacity of the vaccine will be reduced or negated by freezing or high temperature and if the directions for use and precautionary measures are not followed as directed, excessive morbidity and mortality may occur. Consult a poultry veterinarian or a diagnostic laboratory for proper diagnosis. Inform the veterinarian or the diagnostic laboratory that the flock has been vaccinated.

ACTIVE INGREDIENTS

Eimeria adenoeides and E. meleagritidis.

FOR USE IN ANIMALS ONLY

Manufactured and marketed by:
Ceva Animal Health, LLC | Tel: 913-894-0230
8735 Rosehill Road, Ste. 300 | Lenexa, KS 66215 | www.ceva.us

Immucox is a registered trademark of Ceva Animal Health Inc.
© 2022 Ceva Animal Health, LLC

