

ULTIFEND®

IBD ND + **RISPENS**

BURSAL DISEASE - MAREK'S DISEASE
NEWCASTLE DISEASE VACCINE
Serotype 3, Live Marek's Disease Vector
& Serotype 1 Live Marek's

APPROVED SPECIES

- Chickens

VACCINE TYPE

- Live HVT vector IBD & ND vaccine in combination with conventional live Rispens Serotype 1 Marek's

VACCINE CONSTRUCT

- Vector: Live Serotype 3 turkey herpes virus (HVT)
- Donor Inserts:
 - IBDV standard strain , VP2 protein
 - NDV, F protein
- Rispens Serotype 1 Marek's

DISEASE PROTECTION

- Infectious Bursal Disease
- Newcastle Disease
- Very Virulent + Marek's Disease

ROUTE OF ADMINISTRATION

- Subcutaneous injection

AGE OF ADMINISTRATION

- SC: 1-day-old chickens or older

PACKAGING (PRODUCT CODE)

- 5 ampules per cane
- 2,000-dose ampule (B37811US)

VACCINE PRESENTATION

- Cell associated
- Frozen in liquid nitrogen

QUALITY

Thoroughly tested for purity, sterility, potency, and safety and compliant with all applicable USDA and Ceva standards.

KEY ADVANTAGES

- One hatchery application to help optimize your Gumboro, Newcastle and Marek's protection
- Effective IBD & ND vaccination in the face of maternal antibodies
- An opportunity to eliminate post vaccination reactions associated with live Newcastle vaccines
- Known safety and efficacy of HVT vector vaccines
- Expected HVT replication and expression of VP2 & F proteins
- Provides Marek's disease protection equal to conventional HVT + Rispens vaccines
- Rispens strain derived from original European Rispens strain, providing protection against very virulent and very virulent + Marek's disease viruses

See reverse side and package label for complete use directions and information.



ULTIFEND®

IBD ND + RISPENS

BURSAL DISEASE - MAREK'S DISEASE NEWCASTLE DISEASE VACCINE Serotype 3, Live Marek's Disease Vector & Serotype 1 Live Marek's

DESCRIPTION

ULTIFEND® IBD ND + Rispens contains a genetically engineered Marek's disease vaccine of serotype 3 (turkey herpesvirus or HVT) expressing infectious bursal disease and Newcastle disease key protective antigens and a serotype 1 (Rispens) Marek's disease vaccine. The bivalent Marek's disease vaccine containing serotypes 1 and 3 is presented in a frozen cell associated form. The cells and virus particles are very fragile and require careful handling to prevent damage or loss of titer in order to achieve optimum efficacy. The vaccine is stored and shipped in frozen form in liquid nitrogen.

STORAGE CONDITIONS

Vaccine ampules: Store in liquid nitrogen

Diluent: Store at room temperature between 68 - 77°F (20 - 25°C)

INDICATIONS

This product has been shown to be effective for the vaccination of healthy chickens 1 day of age or older against standard bursal disease, Newcastle disease and very virulent Marek's disease. The duration of immunity is unknown. For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

Good management practices are recommended to reduce exposure to bursal disease virus, Newcastle disease virus and Marek's disease virus for at least two weeks following vaccination.

PREPARATION AND ADMINISTRATION OF VACCINE

Carefully read the directions before use. The instructions must be completely followed.

1. For subcutaneous injection of day-old chicks: Mix 200 ml of diluent for each 1000 doses of vaccine.

VACCINE PREPARATION

1. Be familiar with all safety and precautionary measures for handling liquid nitrogen to prevent personal injury.
2. Wear gloves, a plastic face shield, and protective goggles before removing vaccine ampules from the liquid nitrogen.
3. Check the container to confirm a sufficient amount of liquid nitrogen is present to keep the vaccine frozen. If thawed, do not use the vaccine. The containers (Dewars) must be checked regularly for liquid nitrogen level and must be refilled as needed.
4. After matching the dose size of the vaccine with the diluent size, quickly remove the exact number of ampules needed.
5. A maximum of 3 ampules should be thawed at one time. After inspecting the diluent and completing all preparations, the vaccine should be quickly removed from the Dewar and placed into a clean 80°F (26.5°C) thaw bath. This step can be repeated

until the appropriate dosage has been reached. Gentle agitation during the thawing process promotes rapid, uniform thawing and evenly distributes the vaccine in the ampule.

6. Immediately after thawing, mix the vaccine with the diluent at room temperature (68°-77°F or 20°-25°C). Gently draw the vaccine from the ampule with an 18-gauge, 1.5 inch needle and slowly mix with the diluent. Rinse the ampule one time with the diluted vaccine.
7. Immediately use the vaccine and mix occasionally to ensure uniform suspension of cells.

SUBCUTANEOUS VACCINATION

1. For subcutaneous injection, sterilize the automatic syringe, needles and other accessory equipment by autoclaving or boiling prior to vaccination.
2. Use a short (3/8 inch or 1/2 inch) 20-gauge needle for vaccination. Subcutaneously inject 0.2 ml into the back of the neck of each chick.

NOTICE

This vaccine has been thoroughly tested for safety, purity, potency and sterility and is in compliance with all applicable USDA and Ceva standards.

CAUTION

1. Read the above directions carefully.
2. Do not vaccinate within 21 days before slaughter.
3. The vaccine contains gentamicin and amphotericin B as preservatives.
4. Inactivate unused contents before disposal.
5. Do not remove vaccine from liquid nitrogen until ready for use.
6. Do not refreeze the vaccine.
7. Use entire contents when first opened.
8. Do not use vaccine that has thawed in the liquid nitrogen container.
9. Do not overdose or over dilute the vaccine.
10. Do not mix with other products, except as specified on this label.
11. In case of human exposure, contact a physician.
12. For use in animals only.
13. The effect of combining this vaccine with other products is not known and could impact vaccine efficacy.

Manufactured by: Ceva Animal Health, LLC

Marketed by: Ceva Animal Health, LLC | Tel: 913-894-0230

8735 Rosehill Road, Ste. 300 | Lenexa, KS 66215 | www.ceva.us

Ultifend registered trademark of Ceva Santé Animale S.A.

