



cevac CIRCOMUNE[®]

• Chicken Infectious Anemia

APPROVED SPECIES

- Chickens (breeders)

VACCINE TYPE

- Live Chicken Anemia virus (CAV)
 - Del-Ros strain

DISEASE PROTECTION

- Chicken Infectious Anemia

VACCINE PRESENTATION

- Ready-to-use liquid presentation

ROUTE OF ADMINISTRATION

- Wing-web

AGE OF ADMINISTRATION

- 9- to 12-week-old chickens

PACKAGING (PRODUCT CODE)

- 10 x 1,000 doses (B333H1US)

QUALITY

- Thoroughly tested for purity, sterility, potency, and safety, and compliant with all applicable USDA and Ceva standards.

Key Advantages

- Del-Ros strain is derived from a natural field strain that is naturally avirulent
- Induces maternal immunity in progeny
- Provides protection against chicken infectious anemia in breeders
- Contains an immunogenic strain of CAV for maximum protection
- Convenient: vaccine does not require addition of diluent

CIRCOMUNE®

CHICKEN ANEMIA VIRUS VACCINE

Live virus

For Wing Web Administration



DESCRIPTION

CIRCOMUNE® is a live chicken infectious anemia virus vaccine, Del-Ros strain, for use in chickens using wing web administration. The vaccine is stable allowing it to be presented in a liquid, ready to use preparation that can be stored at 35°-45°F (2°-7°C).

INDICATIONS

CIRCOMUNE is recommended for use only by wing web administration to healthy, susceptible chickens 9 to 12 weeks of age. Maternal antibodies are transferred to progeny to aid in the prevention of disease caused by chicken infectious anemia virus.

ADMINISTRATION

Shake well before opening. Remove seal and stopper from the vaccine vial. The vaccine is ready to use. Carefully hold each individual bird and spread the wing with the underside facing up. Dip the double-pronged applicator into the vaccine so that the needle grooves fill with liquid to deliver 0.01 mL per bird. Insert the applicator into the webbed portion of the wing avoiding feathers, muscle, bone and large blood vessels. Re-dip the applicator between each vaccination.

WARNING

Use of this vaccine in chickens younger than 3 weeks of age may cause clinical signs of chicken infectious anemia. Avoid spread of vaccine virus to young chickens or unexposed laying chickens. This vaccine is not recommended for chickens older than 12 weeks of age. Controlled studies demonstrated that this vaccine virus did not vertically transmit to progeny. However, it is not recommended to vaccinate chickens in lay to prevent the possibility of the vaccine virus being transmitted to progeny.

PRECAUTIONS

Store this vaccine at 35°-45°F (2°-7°C). Vaccinate only healthy birds. All susceptible chickens should be vaccinated at the same time. Do not dilute the vaccine or administer less than 1 dose per bird. Do not vaccinate within 21 days of slaughter. Burn open vials and unused contents. Contains gentamicin and amphotericin B as preservatives.

QUALITY ASSURANCE

Every serial of this vaccine has been carefully produced and has undergone purity, sterility, potency and safety tests to meet Ceva's quality standards and USDA regulations.



Manufactured by:
Biomune Company

Distributed by:
Ceva Animal Health, LLC
8735 Rosehill Road, Ste. 300
Lenexa, KS 66215

Tel: 913-894-0230
www.ceva.us

All trademarks are owned by Ceva Santé Animale S.A. and its affiliates.